

Effects of dietary thiamin on the physiological status of the grouper *Epinephelus coioides*

J. W. Huang · L. X. Tian · Z. Y. Du ·
H. J. Yang · Y. J. Liu

Received: 18 June 2006 / Accepted: 4 January 2007 / Published online: 22 February 2007
© Springer Science+Business Media B.V. 2007

Abstract This study was conducted to evaluate the effects of dietary thiamin on the physiological status of the juvenile grouper, *Epinephelus coioides*. Graded levels of thiamin (0.08, 0.50, 2.12, 3.15, 4.63, 12.37 mg thiamin kg⁻¹ diet) were fed to grouper juveniles (mean weight: 16.97 ± 0.14 g) for 10 weeks. Although fish fed the thiamin-deficient (TD) diet showed no obvious symptoms of thiamin deficiency or increased mortality, those fed the lowest doses of thiamin (0.08 and 0.50 mg thiamin kg⁻¹ diet) had significantly decreased transketolase activity in the liver. In addition, the level of liver thiobarbituric acid reactive substances in fish fed the TD diet was 33–67% higher than that in fish with the thiamin-supplemented diet. There were no significant differences in superoxide dismutase activity between the different groups of fish.

Keywords *Epinephelus coioides* · Grouper · Superoxide dismutase (SOD) · Thiamin · Thiobarbituric acid reactive substances (TBARs) · Transketolase

Introduction

The water-soluble vitamin, thiamin, is an essential nutrient in the diet of humans and other animals, and the detrimental effects of thiamin deficiency originating from alcoholism and brain diseases have been reported under in vitro and in vivo conditions (Ba et al. 2005; Hazell et al. 2003; Lonsdale 2006; Martin et al. 2003). In addition, the requirements for thiamin and the symptoms of thiamin deficiency have been determined in a number of fish species (Amcoff et al. 2002; Hashimoto et al. 1970; Lehmitz and Spannhof 1977; Masumoto et al. 1987; Morito et al. 1986; Murai and Andrews 1978). Severe thiamin deficiency in mammals is often associated with neurological disturbances (Ciccia and Langlais 2000; Mulholland 2006; Nakagawasai et al. 2000; Pannunzio et al. 2000) and is strongly related to impaired carbohydrate metabolism in the form of thiamin pyrophosphate (TPP) (Martin et al. 2003).

TPP is an active form of thiamin, and it acts as an important coenzyme for transketolase and several other enzymes involved in the conversion of carbohydrate and fat into energy, including pyruvate dehydrogenase and α -ketoglutarate dehydrogenase, which are important in carbohydrate metabolism and intracellular redox homeostasis (Blair et al. 1999; Shangari et al. 2003). Transketolase activity is very sensitive to thiamin

J. W. Huang · L. X. Tian · Z. Y. Du · H. J. Yang ·
Y. J. Liu (✉)
Institute of Aquatic Economic Animals, School of
Life Sciences, Sun Yat-sen University, 135 Xingang
West Street, Haizhu District, Guangzhou 510275
Guangdong, P.R. China
e-mail: Fishnutri@163.com

status and has been shown to be the rate-limiting enzyme in the nonoxidative portion of the pentose phosphate pathway (Berthon et al. 1992). A thiamin deficiency-induced impairment of carbohydrate metabolism can lead to cellular oxidative stress (Calingasan et al. 1999; Langlais et al. 1997; Todd and Butterworth 1999), while a number of studies in the rat have suggested that thiamin has antioxidative properties (Langlais et al. 1997; Lukienko et al. 2001).

Wilson (1994) pointed out that fish cannot utilize carbohydrate efficiently; this is in contrast to the situation in mammals where carbohydrate is one of the main sources of fuel/energy. This has raised the question of whether there would be a difference between the rat and fish in terms of their degree of dependence on thiamin. However, to date very little physiological research has focused on the role of thiamin in fish. In Asia, the grouper, *Epinephelus coioides*, is considered to be an important commercial maricultured carnivorous fish (Boonyaratpalin 1997), but very little information has been published on the role of the B-family vitamins in this fish; the exception being pyridoxine (Huang et al. 2005). Our previous study (Huang et al. 2005) reported that symptoms of pyridoxine deficiency in the grouper resemble those reported in mammals (Baxter 2003). Because pyridoxine and thiamin are tightly correlated to protein and carbohydrate metabolisms, the aim of this study was to determine some of the physiological and biochemical alterations that occur in grouper as a result of thiamin deficiency, including changes in the levels of transketolase, superoxide dismutase (SOD), and oxidative stress, in order to compare the differences in thiamin-related metabolic processes between fish and mammals.

Materials and methods

Diet preparation

The formulation of the basic purified diet (45.8% protein and 14.18 kJ g⁻¹ gross energy) – excluding the thiamin components – is shown in Table 1. The vitamin-free casein was supplied by Sigma-Aldrich (product no. C3400; St. Louis, Mo.); all

other ingredients were obtained from companies located in the People's Republic of China. Thiamin hydrochloride (Roche) was added to the diets at the expense of cellulose to provide concentrations of 0, 1, 2.5, 5, 10, and 20 mg kg⁻¹ dry diet. The diets were analyzed for thiamin by the fluorometric method (AOAC 1990) and found to contain 0.08, 0.50, 2.12, 3.15, 4.63, and 12.37 mg kg⁻¹ dry diet, respectively. The diets were prepared by first mixing the dry ingredients with oil and water, followed by pelleting the wet mixture into 2.5-mm pellets and air-drying in an air-conditioned room for two nights (2000–0800 hours). The air-dried pellets were stored at –20°C until use.

Experimental procedure

Grouper juveniles were obtained from a local fish fry dealer. Upon arrival, they were acclimatized to laboratory conditions for 4 weeks. The fish were fed the thiamin-deficient (TD) diet for 2 weeks prior to the trial to deplete the body of all stored thiamin; subsequently, the experimental diets were fed to triplicate groups of 20 fish, with an individual initial weight 16.97 ± 0.14 g. The experiments were conducted in flow-through seawater tanks (300 l), provided with continuous oxygen by an air stone connected to the air pump. The water temperature was maintained between 27.3°C and 31.9°C, dissolved oxygen content was ≥6 mg l⁻¹, and the salinity was 28–30‰. A 12:12-h photoperiod was maintained (light: 0700–1900 hours). The trial lasted for 10 weeks. For the duration of the trial, any abnormal behavior of the fish was observed and recorded daily.

Assays for biochemical enzymes

At the end of the experiment, the fish were first subjected to a 24-h fasting period, following which three fish were sacrificed and their livers removed for the enzyme assays. Transketolase (EC 2.2.2.1) activity was measured using the orcinol reaction determination of the sedoheptulose (colorimetric assays; Horecker 1957; Warnock 1970). Liver samples were homogenized and the supernatants incubated with D-ribose-5-phosphate (0.012 M) in a 37°C water bath for 1 h. A trichloroacetic acid

Table 1 Composition of the basal diet, with the exception of thiamin

Ingredients	(g kg ⁻¹ dry weight)
Vitamin-free casein	300
Gelatin	75
Corn starch	200
Fish oil	45
Corn oil	45
Amino acid mixture ^a	120
Choline chloride	5
Ascorbic acid phosphate ester	5
Vitamin mixture ^b	20
Mineral mixture ^c	40
α -Cellulose	145
Protein	458
Carbohydrate	197
Lipid	86
Energy (kJ g ⁻¹)	14.18

^a The composition of the amino acid mixture in grams per kilogram dry diet is: leucine, 15.48; valine, 11.5; phenylalanine, 10.24; tyrosine, 9.68; isoleucine, 9.18; threonine, 8.97; aspartic acid, 7.56; glutamic acid, 7.56; serine, 7.51; proline, 7.46; arginine-HCl, 7.06; lysine-HCl, 5.55; histidine, 4.74; cysteine, 2.57; methionine, 2.47; tryptophan, 2.47 (modified from Shaik Mohamed 2003)

^b The composition of the vitamin mixture in terms of milligrams per gram mixture is: riboflavin, 6.67; calcium pantothenate, 9.33; nicotinic acid, 26.67; pyridoxine hydrochloride, 2.5; folic acid, 0.5; inositol, 133.33; menadione, 1.33; alpha-tocopheryl acetate, 13.33; retinol acetate, 0.403; cholecalciferol, 0.0025; biotin, 2; vitamin B₁₂, 0.003. All ingredients were diluted with alpha-cellulose to 1 g

^c The composition of the mineral mixture in terms of milligrams per gram mixture is: calcium lactate, 333.3; KCl, 233.3; Ca(H₂PO₄)₂·H₂O, 133.3; MgSO₄·7H₂O, 133.3; NaH₂PO₄·H₂O, 83.3; FeSO₄·H₂O, 33.3; ZnSO₄·7H₂O, 20; MnSO₄·4H₂O, 6.7; KI, 6.7; AlCl₃·6H₂O, 6.7; CuSO₄·5H₂O, 10

solution (15%) was then added to terminate the enzymatic reaction. A fixed volume of the solution was centrifuged at 10000 rpm for 10 min, following which ferric chloride-orninol reagent was added to the supernatant and the mixture placed in a boiling water bath for 40 min. The optical densities were then measured at 580 and 670 nm using a UV/VIS spectrophotometer. Transketolase activity was calculated using the method of Warnock (1970), and the activity of this enzyme was expressed as micromoles sedoheptulose per gram wet tissue. Liver SOD activity was determined using a commercial kit (Jiancheng Bioengineering Institute, Nanjing,

P.R. China). The results are expressed as the percentage inhibition of color formation, with high values indicating high SOD activity.

Assay for liver thiobarbituric acid reactive substances (TBARs)

The susceptibility of liver to induced lipid peroxidation (LPO) was assayed by the TBARs test using a commercial kit (Jiancheng Bioengineering Institute). The results are expressed as nanomoles of malondialdehyde (MDA) per milligram protein. Protein measurements were carried out by the Bradford method using a commercial kit (Beyotime Biotechnology, Jiangsu, P.R. China).

Statistical analysis

All data were subjected to one-way analysis of variance (ANOVA) using the SPSS ver. 12.0 statistical software (SPSS, Chicago, Ill.). Significant differences between means were analyzed using Duncan's new multiple range test at a 95% interval of confidence ($P < 0.05$).

Results

During the 10-week experimental period, no obvious signs of thiamin deficiency were observed. As shown in Fig. 1, there was no significant difference in terms of the survival of *E. coioides* (90.0–98.3%; $P > 0.05$) between the groups fed the different diets.




Fig. 1 Survival of *Epinephelus coioides* fed diets containing different levels of thiamin for 10 weeks. Values are shown as the mean \pm standard deviation (SD) ($n = 3$)




Fig. 2 Liver transketolase activity of *E. coioides* fed diets containing different levels of thiamin for 10 weeks. Values are shown as the mean \pm SD ($n = 3$). Statistical group differences ($P < 0.05$) are marked with *different letters*

The activity of liver transketolase in *E. coioides* was significantly affected by dietary thiamin level only between the lowest doses (0.08 and 0.50 mg kg⁻¹ diet) and the maximum dose (12.37 mg kg⁻¹ diet) ($P < 0.05$). There was no significant difference when the diet contained thiamin at levels between 0.08 to 4.63 mg kg⁻¹ (Fig. 2).

Levels of TBARs in the liver decreased from 8.82 to 5.29 nmol MDA mg⁻¹ protein as the dietary thiamin increased from 0.08 to 12.37 mg kg⁻¹ diet (Fig. 3a). TD fish had 33–67% higher TBARs levels




Fig. 3 Content of liver thiobarbituric acid reactive substances (TBARs) (a) and superoxidase dismutase (SOD) activity (b) of *E. coioides* fed diets containing different levels of thiamin for 10 weeks. Values are shown as the mean \pm SD ($n = 3$). Statistical group differences ($P < 0.05$) are marked with *different letters*

than the fish fed the thiamin-supplemented diets; significant difference was detected between the TD fish and those fed 3.15 to 12.37 mg kg⁻¹ of thiamin ($P < 0.05$). There was no significant difference among all the thiamin-supplemented groups ($P > 0.05$). Figure 3b shows that thiamin supplemented to the diet had no effect on liver SOD activity ($P > 0.05$).

Discussion

There were no obvious signs of thiamin deficiency in the 10-week experiment reported here. A similar result was also reported for turbot fed a TD diet for 16 weeks (Cowe et al. 1975). Murai and Andrews (1978) reported the absence of neurological symptoms in channel catfish fed a TD diet for 20 weeks, although other signs of thiamin deficiency, such as anorexia, dark coloration of the skin, extremely poor growth, and increased mortality rates, did appear. Neurological symptoms, such as ataxia, were reported on rainbow trout fed with a TD diet for 30 weeks (Masumoto et al. 1987). On the basis of our results and those reported earlier, it would appear that TD symptoms, especially neurological symptoms, are not easy to induce in fish. When these results on thiamin deficiency in fish are compared to the thiamin deficiency symptoms reported in rats, it is interesting to note that fish may be much less susceptible to thiamin deficiency (at least in terms of the more obvious symptoms) than rats. For example, neurological symptoms appeared in rats fed a TD diet for only about 4 weeks (Ciccia and Langlais 2000); furthermore, when treated with a thiamin antagonist, pyriethamine, rats showed gross neurological symptoms after only 14 days of treatment (Gibson et al. 1984). Thiamin plays an important role in carbohydrate metabolism, and the energy derived from the oxidation of glucose is highly dependent upon the availability of TPP (Lonsdale 2006). The ingestion of carbohydrate automatically increases the need for dietary thiamin (Elmadfa et al. 2001; Lonsdale 2006). Compared to mammals, fish utilize carbohydrate relatively poorly (Wilson 1994). Therefore, fish may need less thiamin and may be less susceptible to thiamin deficiency than mammals.

Alterations in the cellular carbohydrate metabolism has been observed to be a common cause of neurological lesions in the TD rat brain (Singleton and Martin 2001). However, fish – especially carnivorous fish – have a remarkable capacity to utilize amino acids both as a metabolic fuel and as precursors for protein, lipid, and carbohydrate synthesis (Wood 1993). In other words, carbohydrate is not the most important source of energy in carnivorous fish (Wilson 1994), and once carbohydrate metabolism is impaired under extreme conditions (such as in thiamin deficiency), fish can still live for a relatively long time. Taking into account that our previous study (Huang et al. 2005) found that grouper fish were very susceptible to pyridoxine deficiency, we suggest that our results emphasize the fact that carbohydrate is relatively less important than protein in carnivorous fish.

Although no clear-cut signs of thiamin deficiency were observed in the present study, liver transketolase activity had been affected by thiamin deficiency. Transketolase activity is regarded as being the most sensitive indicator of thiamin status, and a substantial decline in transketolase activity resulting from thiamin deficiency has been found to occur in the brain of mammals prior to the onset of severe brain damage (Giguère and Butterworth 1987). The decrease in transketolase activity in the present study was similar to that observed in the rat (Rains et al. 1997) and other aquatic animals, such as turbot (Cowey et al. 1975), abalone (Zhu et al. 2002) and shrimp (Chen et al. 1994). Sheu et al. (1996) suggested that the decrease in transketolase activity is due to a decrease in the enzyme protein. This loss of transketolase protein was demonstrated to be due to a decrease in its synthesis rate and mRNA steady state levels but not to an increase in the degradation rate, as reported by Pekovich et al. (1998).

NADPH can be produced to combat oxygen radicals in the transketolase-mediated pentose phosphate pathway (Berthon et al. 1992). As such, the disruption in the metabolism of transketolase can destroy the homeostasis of cellular redox potential (Sheu et al. 1996) and can further lead to oxidative stress (Langlais et al. 1997). Oxidative stress is considered to be one of the major factors

underlying the deleterious effects of thiamin deficiency. The present results indicate that the highest TBAR values occurred in fish fed the TD diet (see Fig. 3a). Similarly, high levels of free radicals or lipid peroxidation (LPO) products were found in TD rats (Calingasan et al. 1999; Langlais et al. 1997). However, the addition of thiamin to the diet at concentrations of 10^{-4} – 10^{-6} M can inhibit LPO in rat liver microsomes, as reported by Lukienko et al. (2001), who subsequently suggested that the antioxidant effects of thiamin result from thiamin being able to interact directly with free radicals and hydroperoxides by successive transfer ($2H + 2e^-$) from the NH_2 group of the pyrimidine ring to radicals. However, whether the grouper was fed with a TD diet or not, we found no significant difference in the activity of liver SOD, which is a free radical scavenging enzyme. We suggest the possibility that SOD is not a main factor affecting oxidative stress in grouper, but further research on the antioxidant effects of thiamin need to be carried out.

Acknowledgements This project was supported by the Key Technologies R&D Program during the tenth 5-year plan, China (grant no. 2001DA505D/06). The authors are grateful to Dr. Wei Zhu (Ocean University of China) for providing some references.

References

- Amcoff P, Elofsson UOE, Börjeson H et al (2002) Alterations of dopaminergic and serotonergic activity in the brain of sea-run Baltic salmon suffering a thiamine deficiency-related disorder. *J Fish Biol* 60:1407–1416
- AOAC (Association of Official Analytical Chemists) (1990) Thiamin (vitamin B₁) in foods. Fluorometric method. *Official Methods of Analysis*, no. 942.23, 15th edn. AOAC, Arlington, Va.
- Ba A, N'Douba V, D'Almeida MA et al (2005) Effects of maternal thiamine deficiencies on the pyramidal and granule cells of the hippocampus of rat pups. *Acta Neurobiol Exp (Wars)* 65:387–398
- Baxter P (2003) Pyridoxine-dependent seizures: a clinical and biochemical conundrum. *Biochim Biophys Acta* 1647:36–41
- Berthon HA, Kuchel PW, Nixon PF (1992) High control coefficient of transketolase in the nonoxidative pentose phosphate pathway of human erythrocytes: NMR, antibody, and computer simulation studies. *Biochemistry* 31:12792–12798
- Blair PV, Kobayashi R, Edwards III HM et al (1999) Dietary thiamin level influences levels of its diphosphate

- form and thiamin-dependent enzymic activities of rat liver. *J Nutr* 129:641–648
- Boonyaratpalin M (1997) Nutrient requirements of marine food fish cultured in Southeast Asia. *Aquaculture* 151:283–313
- Calingasan NY, Chun WJ, Park LC et al (1999) Oxidative stress is associated with region-specific neuronal death during thiamine deficiency. *J Neuropathol Exp Neurol* 58:946–958
- Chen HY, Wu FC, Tang SY (1994) Sensitivity of transketolase to the thiamin status of juvenile marine shrimp (*Penaeus monodon*). *Comp Biochem Phys A* 109:655–659
- Ciccia RM, Langlais PJ (2000) An examination of the synergistic interaction of ethanol and thiamine deficiency in the development of neurological signs and long-term cognitive and memory impairments. *Alcohol Clin Exp Res* 24:622–634
- Cowey CB, Adron JW, Knox D et al. (1975) Studies on the nutrition of marine flatfish. The thiamine requirement of turbot (*Scophthalmus maximus*). *Br J Nutr* 34:383–390
- Elmadfa I, Majchrzak D, Rust P et al. (2001) The thiamine status of adult humans depends on carbohydrate intake. *Int J Vitam Nutr Res* 71:217–221
- Gibson GE, Ksiezak-Reding H, Sheu K-FR et al (1984) Correlation of enzymatic, metabolic, and behavioral deficits in thiamine deficiency and its reversal. *Neurochem Res* 9:803–814
- Giguère J-F, Butterworth RF (1987) Activities of thiamine-dependent enzymes in two experimental models of thiamine deficiency encephalopathy: 3. transketolase. *Neurochem Res* 12:305–310
- Hashimoto Y, Arai S, Nose T (1970) Thiamine deficiency symptoms experimentally induced in the eel. *Nippon Suisan Gakkai Shi* 36:791–797
- Hazell AS, Pannunzio P, Rama Rao KV et al (2003) Thiamine deficiency results in downregulation of the GLAST glutamate transporter in cultured astrocytes. *Glia* 43:175–184
- Horecker BL (1957) The orcinol reaction for mixtures of pentose and heptulose. In: Colowick SP, Kaplan NO (eds) *Methods of enzymology III*. Academic Press, New York, pp 105–107
- Huang JW, Tian LX, Du ZY et al (2005) Pyridoxine deficiency of grouper, *Epinephelus coioides*: physiological and biochemical alteration. *Fish Physiol Biochem* 31:331–337
- Langlais PJ, Anderson G, Gu SX et al (1997) Increased cerebral free radical production during thiamine deficiency. *Metab Brain Dis* 12:137–143
- Lehmitz R, Spannhof L (1977) Transketolase activity in thiamine deficiency of the kidney in rainbow trout (*Salmo gairdneri*) following continuous feeding with raw herring. *Arch Tierernahr* 27:287–295
- Lonsdale D (2006) A review of the biochemistry, metabolism and clinical benefits of thiamin(e) and its derivatives. *Evid Based Complement Alternat Med* 3:49–59
- Lukienko PI, Mel'nichenko NG, Zverinskii IV et al (2001) Antioxidant properties of thiamine. *Bull Exp Biol Med* 130:874–876
- Martin PR, Singleton CK, Hiller-Sturmhöfel S (2003) The role of thiamine deficiency in alcoholic brain disease. *Alcohol Res Health* 27:134–142
- Masumoto T, Hardy RW, Casillas E (1987) Comparison of transketolase activity and thiamin pyrophosphate levels in erythrocytes and liver of rainbow trout (*Salmo gairdneri*) as indicators of thiamine status. *J Nutr* 117:1422–1426
- Morito CLH, Conrad DH, Hilton JW (1986) The thiamine deficiency signs and requirement of rainbow trout (*Salmo gairdneri*, Richardson). *Fish Physiol Biochem* 1:93–104
- Mulholland PJ (2006) Susceptibility of the cerebellum to thiamine deficiency. *Cerebellum* 5:55–63
- Murai T, Andrews JW (1978) Thiamin requirement of channel catfish fingerlings. *J Nutr* 108:176–180
- Nakagawasai O, Tadano T, Nijjima F et al (2000) Immunohistochemical estimation of rat brain somatostatin on avoidance learning impairment induced by thiamine deficiency. *Brain Res Bull* 51:47–55
- Pannunzio P, Hazell AS, Pannunzio M et al (2000) Thiamine deficiency results in metabolic acidosis and energy failure in cerebellar granule cells: an in vitro model for the study of cell death mechanisms in Wernicke's encephalopathy. *J Neurosci Res* 62:286–292
- Pekovich SR, Martin PR, Singleton CK (1998) Thiamine deficiency decreases steady-state transketolase and pyruvate dehydrogenase but not α -ketoglutarate dehydrogenase mRNA levels in three human cell types. *J Nutr* 128:683–687
- Rains TM, Emmert JL, Baker DH et al (1997) Minimum thiamin requirement of weanling Sprague-Dawley outbred rats. *J Nutr* 127:167–170
- Shaik Mohamed J, Sivaram V, Christopher Roy TS et al (2003) Dietary vitamin A requirement of juvenile greasy grouper (*Epinephelus tauvina*). *Aquaculture* 219:693–701
- Shangari N, Bruce WR, Poon R et al (2003) Toxicity of glyoxals-role of oxidative stress, metabolic detoxification and thiamine deficiency. *Biochem Soc Trans* 31:1390–1393
- Sheu K-FR, Calingasan NY, Diemel GA et al (1996). Regional reductions of transketolase in thiamine-deficient rat brain. *J Neurochem* 67:684–691
- Singleton CK, Martin PR (2001) Molecular mechanisms of thiamine utilization. *Curr Mol Med* 1:197–207
- Todd KG, Butterworth RF (1999) Early microglial response in experimental thiamine deficiency: an immunohistochemical analysis. *Glia* 25:190–198
- Warnock LG (1970) A new approach to erythrocyte transketolase measurement. *J Nutr* 100:1057–1062
- Wilson RP (1994) Utilization of dietary carbohydrate by fish. *Aquaculture* 124:67–80
- Wood CM (1993) Ammonia and urea metabolism and excretion. In: Evans DH (ed) *The physiology of fishes*. CRC Press, Boca Raton, pp 384–385
- Zhu W, Mai KS, Wu GT (2002) Thiamin requirement of juvenile abalone, *Haliotis discus hannai* Ino. *Aquaculture* 207:331–343