

## Salidroside inhibits H<sub>2</sub>O<sub>2</sub>-induced apoptosis in PC12 cells by preventing cytochrome *c* release and inactivating of caspase cascade

Lei Cai<sup>1</sup>, Hua Wang<sup>2</sup>, Qin Li<sup>2</sup>, Yunfei Qian<sup>1</sup>, and Wenbing Yao<sup>1\*</sup>

<sup>1</sup> School of Life Science and Technology, China Pharmaceutical University, Nanjing 210009, China

<sup>2</sup> Yangtze River Pharmacy Group, Guangzhou Hairui Pharmaceutical Company, Guangzhou 510663, China

**We used a rat pheochromocytoma (PC12) cell line to study the effects of salidroside on hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>)-induced apoptosis. In PC12 cells, H<sub>2</sub>O<sub>2</sub>-induced apoptosis was accompanied by the down-regulation of Bcl-2, the up-regulation of Bax, the release of mitochondrial cytochrome *c* to cytosol, and the activation of caspase-3, -8 and -9. However, salidroside suppressed the down-regulation of Bcl-2, the up-regulation of Bax and the release of mitochondrial cytochrome *c* to cytosol. Moreover, salidroside attenuated caspase-3, -8 and -9 activation, and eventually protected cells against H<sub>2</sub>O<sub>2</sub>-induced apoptosis. Taken together, these results suggest that treatment of PC12 cells with salidroside can block H<sub>2</sub>O<sub>2</sub>-induced apoptosis by regulating Bcl-2 family members and by suppressing cytochrome *c* release and caspase cascade activation.**

**Keywords** salidroside; hydrogen peroxide; apoptosis; PC12 cells

Alzheimer's disease (AD) is a multifaceted neurodegenerative disorder characterized by the progressive deterioration of cognition and memory in association with widespread neuronal loss and the deposit of senile plaques. To date, the cause and the mechanism by which neurons die as a result of AD still remain unclear, yet several lines of evidence support the involvement of apoptosis. Studies on post-mortem tissues have provided direct morphological and biochemical evidence that some neurons in the brains of AD patients degenerate via an apoptotic mechanism

relating to the presence of DNA damage, nuclear apoptotic bodies, and other markers of apoptosis [1,2]. These results suggest therapeutic strategies aimed at preventing and delaying apoptosis might be a reasonable choice for the treatment of the disease.

Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), a major source of reactive oxygen species, destroys neurons by inducing apoptosis, which has implications for several biological and pathological processes, including AD. H<sub>2</sub>O<sub>2</sub> has been used in many studies to trigger cell apoptosis [3,4]. Therefore, we used H<sub>2</sub>O<sub>2</sub> to induce apoptosis in PC12 cells in present study.

Considerable efforts have been made to find natural substances with neuroprotective potential, and attention has been focused particularly on Chinese medicinal plants with nootropic effects. Some plants have been used for thousands of years in China to improve cognition or as anti-aging remedies. In our search for new ingredients from traditional Chinese medicinal herbs, salidroside, a phenolic glycoside involved in cell anti-apoptosis processes [5], was isolated from the rhizome of *Rhodiola rosea* L. (Crassulaceae). However, the neuroprotective role of salidroside is unclear. The present study's aim was to explore whether salidroside could inhibit H<sub>2</sub>O<sub>2</sub>-induced toxicity in PC12 cells and the possible mechanism.

## Materials and Methods

### Materials

Salidroside was purchased from National Institute for the Control of Pharmaceutical and Biological Products (Beijing, China). MTT, fluorescent DNA-binding dye Hoechst 33258, and propidium iodide were purchased from Sigma-Aldrich (St. Louis, USA). Dulbecco's Modified Eagle's Medium (DMEM) and fetal bovine serum were obtained from Gibco Life Technologies (Grand Island, USA). Lactic dehydrogenase (LDH) activity assay kit was obtained from Jiancheng Institute of Biotechnology (Nanjing, China).

Received: March 11, 2008 Accepted: June 17, 2008

This work was supported by grants from the Teaching and Research Award Program for Outstanding Young Teachers (No. 2002-383), the Program for New Century Excellent Talents in University (No. NCET-04-0506) and the Traditional Chinese Medicine Research Foundation of Science and Technology (No. 04-05ZP33)

\*Corresponding author: Tel, 86-25-83271218; Fax, 86-25-83271218; E-mail, wbyao@cpu.edu.cn

Antibody of cytochrome *c* was purchased from Santa Cruz Biotechnology (Santa Cruz, USA). DNA extraction kit and caspase-3, -8 and -9 activity kits were from Beyotime Institute of Biotechnology (Nantong, China). All other chemicals and reagents were of analytical grade.

#### Cell culture and treatment and analysis of cell viability

Cells were cultured and treated as described by Qian *et al* [6]. Briefly, PC12 cells were maintained in DMEM supplemented with heat-inactivated 10% fetal bovine serum, 100 U/ml penicillin, 100 µg/ml streptomycin in a water-saturated 5% CO<sub>2</sub> atmosphere at 37 °C. Experiments were carried out 48 h after cells were seeded into 24-well plates. To produce oxidative stress, H<sub>2</sub>O<sub>2</sub> was freshly prepared from 30% stock solution prior to each experiment, and after 12 h exposure, the level of cellular MTT was quantified as described by Chen *et al* [7]. Cells in 24-well plates were briefly rinsed with phosphate-buffered saline (PBS), and 0.5 mg/ml MTT was added to each well. The microplate was incubated at 37 °C for an additional 4 h. At the end of the incubation, the medium with MTT was removed and 500 µl dimethyl sulfoxide was added to each well. The plate was shaken on a microplate shaker to dissolve the blue MTT-formazan. The absorbance was read at 570 nm on a microplate reader. When the effects of salidroside on the PC12 cells were studied, different concentrations of salidroside were added simultaneous to the medium just before the H<sub>2</sub>O<sub>2</sub> was added.

#### Measurement LDH release

LDH release was measured according to the method of Kruman *et al* [8]. Cells were cultured in 24-well culture plates at a density of 1×10<sup>4</sup> cells/well for LDH assay. After 12 h exposure to H<sub>2</sub>O<sub>2</sub>, LDH activities in the medium were measured using an assay kit according to the manufacturer's instructions.

#### Hoechst staining

To quantify and assess nuclear morphology, PC12 cells were fixed for 10 min with 4% paraformaldehyde in PBS. The cells were then stained for 10 min with 10 µg/ml fluorescent DNA-binding dye Hoechst 33258 to reveal nuclear condensation [9]. Hoechst-stained cells were visualized and photographed under a Leica DMIL microscope (Nussloch, Germany).

#### Analysis of DNA fragmentation

Fragmented DNA was isolated using a DNA extraction kit according to the manufacturer's instructions. The elutriants containing DNA pellets were electrophoresed on a 1.8%

agarose gel at 80 V for 1.5 h. The gel was examined and photographed using an ultraviolet gel documentation system.

#### Flow cytometric analysis of DNA content

DNA content was measured according to the methods of Weinmann *et al* [10]. Briefly, cells were collected and washed with ice-cold PBS and fixed with 70% ethanol. The fixed cells were harvested by centrifugation at 1000 *g* for 5 min; dissolved in 100 µl PBS containing 50 mg/ml RNase A, 50 mg/ml propidium iodide, 0.1% Triton X-100 and 0.1 mM EDTA (pH 7.4); and then incubated at 37 °C for 30 min. The fluorescence of cell was measured by flow cytometer (FACSCalibur; Becton Dickinson, San Jose, USA).

#### Reverse transcription-polymerase chain reaction (RT-PCR) analysis

Total RNA was extracted from PC12 cells, and the potential residual genomic DNA was eliminated with RNase-free-DNase I for 30 min at 37 °C. First-strand complementary DNA was synthesized as follows: 1 h at 42 °C with 100 U Moloney murine leukemia virus reverse transcriptase (Promega, Madison, USA), 15 U ribonuclease inhibitor (Promega), 500 µM dNTP, 0.5 µg oligo(dT)<sub>18</sub> and 2 µg total RNA in a final volume 25 µl, and then 5 min at 95 °C. For PCR amplification, the specific primers included the control glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*, 213 bp): 5'-ATTCAACGGCACAGTCAAGG-3' (forward) and 3'-AGTAGAGGCGGGGAAGACG-5' (reverse); *Bcl-2* (303 bp): 5'-GATGACTTCTCTCGTCGCTA-3' (forward) and 3'-TACGGAAACACCTTGATATA-5' (reverse); *Bax* (331 bp): 5'-GAACTGGACAATAATATGGA-3' (forward) and 3'-TCACTGGTAGAAACACCGAC-5' (reverse). The PCR mixture contained 0.8 pM forward and reverse primers of the *Bax* or *Bcl-2*, 0.4 pM forward and reverse primers of the *GAPDH*, 2.0 mM MgCl<sub>2</sub>, 200 µM deoxyribonucleotide triphosphate, and 1.5 U Taq DNA polymerase. The PCR procedure was performed at 94 °C for 5 min, followed by 28 cycles at 94 °C for 1 min, 51 °C for 30 s, 72 °C for 45 s and extension at 72 °C for 10 min. Next, 10 µl PCR products was mixed with 2 µl loading solution, and electrophoresed on agarose-ethidium bromide gel at 100 V for 1 h. The gels were examined and analyzed by an ultraviolet gel documentation system.

#### Analysis of caspase-3, -8, and -9 activities

Caspase-3, -8, and -9 activities were measured using assay kits according to the manufacturer's instructions. Supernatant was mixed with buffer containing the recognition sequence for caspase attached to p-nitroanilide. The

absorbance of p-nitroanilide was determined at 405 nm. The caspase activities were expressed as percentage compared to control.

### Western blot analysis of cytochrome c

Cell lysates were prepared as described by Jia *et al* [11]. To ensure equal loading of the protein samples, protein concentrations of the cell lysates were determined by Bradford assay. Equal amounts of protein (30 µg in total) were separated by 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis and transferred to a nitrocellulose membrane. The membrane was blocked with 5% skim milk in 1×Tris-buffered saline containing 0.05% Tween-20 (TBST) for 1 h. After blocking, the membrane was incubated with 1% skim milk in TBST, containing the primary mouse monoclonal antibody against cytochrome c (1:500) overnight. The membranes were then washed three times with 1×TBST and then incubated with 1% skim milk in TBST, containing a peroxidase-conjugated goat anti-mouse immunoglobulin G secondary antibody (1:5000) (ZSGB-BIO, Beijing, China). The detection of protein bands was performed using the 3,3'-diaminobenzidine tetrahydrochloride substrate kit (ZSGB-BIO).

### Statistical analysis

All experiments were performed in triplicate. Data are presented as mean±SD. The Duncan test and one-way ANOVA were used for multiple comparisons using SPSS 12.0 software (SPSS, Chicago, USA).

## Results

### Inhibition of H<sub>2</sub>O<sub>2</sub>-induced cytotoxicity by salidroside

In PC12 cells, the protective effect on H<sub>2</sub>O<sub>2</sub>-induced cytotoxicity was assessed by MTT assay after 12 h incubation. As shown in **Table 1**, when the cells were

**Table 1** Protective effect of salidroside on PC12 cells with H<sub>2</sub>O<sub>2</sub>-induced injury assessed by MTT assay

Group	A570	Survival (%)
Normal cells	1.82±0.00	100.0
Injured cells	1.37±0.07 <sup>##</sup>	75.3
Injured cells+100 µM sal	1.80±0.03 <sup>**</sup>	98.9
Injured cells+10 µM sal	1.54±0.17 <sup>**</sup>	84.6
Injured cells+1 µM sal	1.37±0.03	75.3

Data are represented as mean±SD. <sup>\*\*</sup>*P*<0.01 versus injured cells group; <sup>##</sup>*P*<0.01 versus control group. sal, salidroside.

pre-incubated with salidroside (10 and 100 µM), H<sub>2</sub>O<sub>2</sub>-induced cell toxicity was significantly reduced in comparison with the control. Necrosis results in a disruption of the cytoplasmic membrane, and the necrotic cells release cytoplasmic LDH and other cytotoxic substances into the medium. We therefore examined the existence of LDH in the cells' culture medium. The LDH index was significantly reduced at doses of 10 and 100 µM in comparison with the control (**Table 2**). The results of MTT and LDH assays showed that salidroside could have a protective effect against H<sub>2</sub>O<sub>2</sub>-induced cytotoxicity.

**Table 2** Inhibitory effect of salidroside on LDH release of PC12 cells under H<sub>2</sub>O<sub>2</sub> treatment

Group	LDH activity (U/ml)	Increment (%)
Normal cells	0.313±0.001	
Injured cells	0.514±0.001 <sup>##</sup>	64.2
Injured cells+100 µM sal	0.377±0.001 <sup>**</sup>	20.4
Injured cells+10 µM sal	0.451±0.002 <sup>**</sup>	44.1
Injured cells+1 µM sal	0.511±0.001	63.3


Data are represented as mean±SD. <sup>\*\*</sup>*P*<0.01 versus injured cells group; <sup>##</sup>*P*<0.01 versus control group. sal, salidroside.

### Salidroside suppresses H<sub>2</sub>O<sub>2</sub>-induced apoptosis


Hoechst 33258 assay revealed the appearance of a collection of multiple chromatin and fragmented apoptotic nuclei after treatment with 0.5 mM H<sub>2</sub>O<sub>2</sub> for 12 h. However, the apoptotic nuclei were significantly reduced when cells were treated with 100 µM salidroside and 0.5 mM H<sub>2</sub>O<sub>2</sub> [**Fig. 1(A)**]. After the PC12 cells were treated with 0.5 mM H<sub>2</sub>O<sub>2</sub> for 12 h, DNA ladder pattern was detected, but salidroside was able to reduce the ladder pattern in a dose-dependent manner [**Fig. 1(B)**]. When the apoptotic cells were analyzed quantitatively by flow cytometry, a significant increase in the apoptotic rate (from 9.78%±0.2% to 32.23%±4.0%) was found after PC12 cells were treated with 0.5 mM H<sub>2</sub>O<sub>2</sub> for 12 h. When PC12 cells were treated with 100 µM salidroside and 0.5 mM H<sub>2</sub>O<sub>2</sub> for 12 h, the percentage of apoptotic cells decreased from 32.23%±4.0% to 18.61%±1.5% [**Fig. 1(C)**].

### Regulation of mRNA expression of Bax or Bcl-2 by salidroside


As shown in **Fig. 2**, after H<sub>2</sub>O<sub>2</sub> treatment for 6 h, mRNA expressions of *Bax* and *Bcl-2* analyzed by RT-PCR analysis showed *Bcl-2* expression began to decrease and *Bax*



**Fig. 1 Salidroside inhibited H<sub>2</sub>O<sub>2</sub>-induced injury in PC12 cells** (A) Morphological analysis of nuclear chromatin in by Hoechst 33258. (i) Control. (ii) After exposure to 0.5 mM H<sub>2</sub>O<sub>2</sub> for 12 h, cells displayed condensed chromatin and apoptotic nuclei. (iii) After exposure to 0.5 mM H<sub>2</sub>O<sub>2</sub> and 100 μM salidroside for 12 h. (B) Agarose gel electrophoresis of DNA fragmentation. Lane 1, DNA ladder marker 1; lane 2, DNA ladder marker 2; lane 3, control; lane 4, 0.5 mM H<sub>2</sub>O<sub>2</sub> for 12 h; lanes 5–7, salidroside (100, 10 and 1 μM, respectively) and 0.5 mM H<sub>2</sub>O<sub>2</sub> treatment for 12 h. (C) Flow cytometric histograms. (i) Control PC12 cells (apoptotic rate is 9.78%±0.2%). (ii) PC12 cells treated with 0.5 mM H<sub>2</sub>O<sub>2</sub> (apoptotic rate is 32.23%±4.0%). (iii) PC12 cells treated simultaneously with 0.5 mM H<sub>2</sub>O<sub>2</sub> and 100 μM salidroside for 12 h (apoptotic rate is 18.61%±1.5%).



**Fig. 2 RT-PCR analysis of mRNA expression of *Bax* and *Bcl-2* in PC12 cells** (A) mRNA expressions of *Bax* (i) and *Bcl-2* (ii) in PC12 cells treated with 0.05 mM H<sub>2</sub>O<sub>2</sub> for indicated time. Lane 1, DNA marker; lane 2, untreated PC12 cells; lanes 3–7, PC12 cells treated with 0.5 mM H<sub>2</sub>O<sub>2</sub> for 3, 6, 9, 12 and 24 h, respectively. (B) After co-treatment with 0.5 mM H<sub>2</sub>O<sub>2</sub> and salidroside (100 and 1 μM, respectively) for 6 h, total RNA was extracted for RT-PCR analysis of *Bax* (i) and *Bcl-2* (ii) expressions. Lane 1, DNA marker; lane 2, untreated cells; lane 3, PC12 cells treated with 0.5 mM H<sub>2</sub>O<sub>2</sub> for 6 h; lanes 4–5, PC12 cells treated with 0.5 mM H<sub>2</sub>O<sub>2</sub> and salidroside (100 and 1 μM, respectively) for 6 h. GAPDH, the internal control.



**Fig. 3 Detection of caspase-3, -8 and -9 activities in PC12 cells** (A) Treatment with 0.05 mM H<sub>2</sub>O<sub>2</sub> causes a time-dependent increase in caspase-3 (i), -8 (ii) and -9 (iii) activities in PC12 cells. Control, untreated cells. (B) PC12 cells were co-treated with 0.05 mM H<sub>2</sub>O<sub>2</sub> and salidroside (100, 10 and 1 μM) for 9 h. The activities of caspase-3 (i), -8 (ii) and -9 (iii) were detected. Control, untreated cells. \**P*<0.05, \*\**P*<0.01 versus control; #*P*<0.05, ##*P*<0.01 versus H<sub>2</sub>O<sub>2</sub> alone.

expression began to increase. The effects of salidroside on mRNA expression were investigated at the same indicated time. The results showed salidroside (100 and 1 μM) significantly raised *Bcl-2* expression and reduced *Bax* in PC12 cells treated with 0.5 mM H<sub>2</sub>O<sub>2</sub> (Fig. 2).

### Salidroside inhibits the activities of caspase-3, -8 and -9

To gain insight into the molecular effector pathway of H<sub>2</sub>O<sub>2</sub>-induced apoptosis, we first examined whether caspases were downstream effectors in H<sub>2</sub>O<sub>2</sub>-mediated apoptosis. H<sub>2</sub>O<sub>2</sub> treatment caused a time-dependent increase in caspase-3, -8 and -9 proteolytic activities. However, when salidroside and H<sub>2</sub>O<sub>2</sub> were added simultaneously to the medium, decreases in the activity of caspase-3, -8 and -9 were detected (Fig. 3).

### Salidroside reduced cytochrome *c* in the cytosol

As indicated in Fig. 4, Western blot analysis revealed that H<sub>2</sub>O<sub>2</sub> treatment caused a progressive accumulation of cytochrome *c* in the cytosol. This was reduced when PC12 cells were treated with salidroside.



**Fig. 4 Total cell lysates extracted from untreated PC12 cells or PC12 cells treated with salidroside and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), and analyzed for cytochrome *c* content by Western blot analysis** (A) Cytochrome *c* expression in cytosol of PC12 cells treated with 0.5 mM H<sub>2</sub>O<sub>2</sub> for indicated times. (B) Effect of 100 μM salidroside on cytochrome *c* release in PC12 cells treated with 0.5 mM H<sub>2</sub>O<sub>2</sub> for 9 h. Control, untreated cells. Model, cells treated with 0.5 mM H<sub>2</sub>O<sub>2</sub> for 9 h.

## Discussion

Recently, researchers have made considerable efforts to search for natural substances with neuroprotective potential, and particular attention has been paid to Chinese

medicinal plants with nootropic effect. The rhizome of *Rhodiola rosea* L. has been used in East Asia as a tonic and anti-aging agent since ancient times. There has been mounting evidence that the extract from the rhizome of *Rhodiola rosea* L. possesses significant neuroprotective activity and antioxidative effects [12,13], although little is known about its pharmacological effects or active ingredients. In a previous study, salidroside was isolated from the rhizome of *Rhodiola rosea* L. and could significantly inhibit O<sub>2</sub><sup>-</sup> or H<sub>2</sub>O<sub>2</sub>-induced neurotoxicity in rat cortical cultures [14]. Earlier results showed that 100 μM salidroside has little effect on PC12 cells, and there was no significant difference compared with control group. The present findings demonstrated that, in PC12 cells, salidroside reduced H<sub>2</sub>O<sub>2</sub>-induced apoptotic death caused by oxidative stress. Treatment with salidroside significantly attenuated increased LDH leakage and decreased viability in differentiated PC12 cells exposed to H<sub>2</sub>O<sub>2</sub>. In these instances, the amount of H<sub>2</sub>O<sub>2</sub> was greater than that of salidroside, and the decrease in cell survival caused by H<sub>2</sub>O<sub>2</sub> was nearly suppressed in the presence of 0.1 mM salidroside. Therefore, we have speculated that antioxidation is just one of salidroside's pathways in this model. Inhibition of relative targets in apoptosis might be a possible mechanism involved in the protective effects of salidroside.

It has been well documented that some pathological neuronal loss in AD occurs through apoptosis. The results of this present study showed that salidroside protected PC12 cells against H<sub>2</sub>O<sub>2</sub>-induced apoptosis. Exposure to 0.5 mM H<sub>2</sub>O<sub>2</sub> induced typical apoptosis in PC12 cells. These results were in accordance with previous studies that found oxidative stress to be a common cause of apoptosis [15, 16]. When cells were pre-incubated with salidroside, H<sub>2</sub>O<sub>2</sub>-induced cell injury was significantly attenuated. For these reasons, salidroside could be a useful neuroprotective agent to ameliorate oxidative stress-induced apoptosis, which may be used in the treatment of AD.

Apoptosis is a type of cell death that represents the culmination of naturally occurring or highly programmed mechanisms. Elucidating the expression patterns of those factors during apoptotic cell death may be critical to our understanding of the underlying mechanisms. Caspase-3 is a key executioner caspase involved in neuronal apoptosis, and its activity is controlled by upstream regulators, such as caspase-8 or caspase-9, which modulate the mitochondria- and death receptor-dependent pathway, respectively [17]. The present study showed that caspase-3 activity was up-regulated in H<sub>2</sub>O<sub>2</sub>-treated cells. We also detected enhanced caspase-9 activity in H<sub>2</sub>O<sub>2</sub>-treated cells

and the release of cytochrome *c* from mitochondria into cytosol. Taken together, these results suggested that H<sub>2</sub>O<sub>2</sub>-induced apoptosis in PC12 cells is associated with the release of cytochrome *c* and the activation of caspases, probably via the mitochondria-mediated apoptosis pathway. We further demonstrated the down-regulation of Bcl-2 or up-regulation of Bax in H<sub>2</sub>O<sub>2</sub>-treated cells. Increased Bax and lowered Bcl-2 expression have been shown to reduce mitochondrial membrane potential and increase reactive oxygen species production in neurons [18], both of which are early events in the process of apoptosis [19]. Our results suggested that the down-regulation of Bcl-2 or up-regulation of Bax alters mitochondrial membrane permeability, triggers mitochondrial cytochrome *c* release to cytosol and activates caspase cascade.

Caspase-8 is a key initiating caspase involved in neuronal apoptosis and that modulates the death receptor-dependent pathway. We detected enhanced caspase-8 activity in H<sub>2</sub>O<sub>2</sub>-treated cells. The results suggested that the death receptor-mediated pathway is involved in H<sub>2</sub>O<sub>2</sub>-induced apoptosis. However, recent studies have suggested that caspase-8 is not always activated early in the context of Fas signaling. In some cells, caspase-9 initiates the processing of caspase-3, which in turn activates caspase-2 and -6. Caspase-6 was found to be required for the activation of downstream caspase-8 [20]. In summary, our study suggested that H<sub>2</sub>O<sub>2</sub>-induced apoptosis in PC12 cells is mediated by at least one pathway through mitochondria that regulates the Bcl-2 family and caspase-3 and -9. However, future studies are required to determine whether the death receptor-mediated pathway is involved in H<sub>2</sub>O<sub>2</sub>-induced apoptosis.

Apoptosis is closely associated with the progression of AD and other neurological diseases. In searching for anti-apoptosis agents, this study examined the possible role of salidroside. Salidroside is an invaluable source for the development of effective neuroprotective agents to protect against apoptosis in PC12 cells in the treatment of age-related neurological diseases.

## References

- 1 Lassmann H, Bancher C, Breitschopf H, Wegiel J, Bobinski M, Jellinger K, Wisniewski HM. Cell death in Alzheimer's disease evaluated by DNA fragmentation *in situ*. *Acta Neuropathol* 1995, 89: 3541
- 2 Smale G, Nichols NR, Brady DR, Finch CE, Horton WE Jr. Evidence for apoptotic cell death in Alzheimer's disease. *Exp Neurol* 1995, 133: 225–230
- 3 Guan S, Bao YM, Jiang B, An LJ. Protective effect of protocatechuic acid from *Alpinia oxyphylla* on hydrogen peroxide-in-

- duced oxidative PC12 cell death. *Eur J Pharmacol* 2006, 538: 73–79
- 4 Tang XQ, Feng JQ, Chen J, Chen PX, Zhi JL, Cui Y, Guo RX *et al.* Protection of oxidative preconditioning against apoptosis induced by H<sub>2</sub>O<sub>2</sub> in PC12 cells: mechanisms via MMP, ROS, and Bcl-2. *Brain Res* 2005, 1057: 57–64
  - 5 Zhang WS, Zhu LQ, Deng RC, Niu FL, Tian R. Effect of salidroside on mitochondrial membrane potential during injury induced by hypoxia/hypoglycemia in cultured SH-SY5Y cells. *Chinese Journal of Pathophysiology* 2004, 20: 1218–1221
  - 6 Qian YF, Wang H, Yao WB, Gao XD. The aqueous extract of the Chinese medicine Danggui-Shaoyao-San inhibits apoptosis in hydrogen peroxide-induced PC12 cells by preventing cytochrome *c* release and inactivating of caspase cascade. *Cell Biol Int* 2008, 32: 304–311
  - 7 Chen P, Li A, Zhang MJ, He ML, Chen Z, Wu XK, Zhao CJ *et al.* Protective effects of a new metalloporphyrin on paraquat-induced oxidative stress and apoptosis in N27 cells. *Acta Biochim Biophys Sin* 2008, 40: 125–132
  - 8 Wang H, Yao WB, Qian YF, Gao XD. Protective effect of salidroside on apoptosis in PC12 cells induced by cyanide and glucose deprivation. *Journal of China Pharmaceutical University* 2007, 38: 273–276
  - 9 Kruman I, Bruce-Keller A, Bredesen D, Waeg G, Mattson MP. Evidence that 4-hydroxynonenal mediates oxidative stress-induced neuronal apoptosis. *J Neurosci* 1997, 17: 5089–5100
  - 10 Weinmann P, Scharffetter-Kochanek K, Forlow SB, Peters T, Walzog B. A role for apoptosis in the control of neutrophil homeostasis in the circulation: insights from CD18-deficient mice. *Blood* 2003, 101: 739–746
  - 11 Jia RR, Gou YL, Ho LS, Ng CP, Tan NH, Chan HC. Anti-apoptotic activity of Bak Foong Pills and its ingredients on 6-hydroxydopamine-induced neurotoxicity in PC12 cells. *Cell Biol Int* 2005, 29: 835–842
  - 12 Xu Q, Guo ZY, Kang JS, Zhu SG, Du KQ, Yang HF. Study on protection of *Rhodiola sachalinensis* AR against free radical injury of the ischemia reperfusion rats. *Journal of Mormon Bethune University of Medical Sciences* 1999, 25: 232–234
  - 13 Lee MW, Lee YA, Park HM, Toh SH, Lee EJ, Jang HD, Kim YH. Antioxidative phenolic compounds from the roots of *Rhodiola sachalinensis* A. Bor. *Arch Pharm Res* 2000, 23: 455–458
  - 14 Li TW, Kong LK, Mu JY, Li XM, Yang HY. The protection of salidroside against O<sub>2</sub><sup>-</sup> or H<sub>2</sub>O<sub>2</sub> damage to rat cortical cultures. *China Academic Journal Electronic Publishing House* 1997, 14: 143–144
  - 15 Whittemore ER, Loo DT, Watt JA, Cotman CW. A detailed analysis of hydrogen peroxide-induced cell death in primary neuronal culture. *Neuroscience* 1995, 67: 921–932
  - 16 Satoh T, Sakai N, Enokido Y, Uchiyama Y, Hatanaka H. Survival factor-insensitive generation of reactive oxygen species induced by serum deprivation in neuronal cells. *Brain Res* 1996, 733: 9–14
  - 17 Kuida K, Zheng TS, Na S, Kuan C, Yang D, Karasuyama H, Ralvic P *et al.* Decreased apoptosis in the brain and premature lethality in CPP32-deficient mice. *Nature* 1996, 384: 368–372
  - 18 Pastorino JG, Chen ST, Tafani M, Snyder JW, Farber JL. The overexpression of Bax produces cell death upon induction of the mitochondrial permeability transition. *J Biol Chem* 1998, 273: 7770–7775
  - 19 Marchetti P, Hirsch T, Zamzami N, Castedo M, Decaudin D, Susin SA, Mace B *et al.* Mitochondrial permeability transition triggers lymphocyte apoptosis. *J Immunol* 1996, 157: 4830–4836
  - 20 Slee EA, Harte MT, Kluck RM, Wolf BB, Casiano CA, Newmeyer DD, Wang HG *et al.* Ordering the cytochrome *c*-initiated caspase cascade: hierarchical activation of caspases-2, -3, -6, -7, -8, and -10 in a caspase-9-dependent manner. *J Cell Biol* 1999, 144: 281–292